

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

Amendment Pursuant to 37 C.F.R. § 1.121

IN THE SPECIFICATION:

IN THE CLAIMS:

The claims set forth below with amendments as indicated will replace all prior versions and listing of claims in the application.

- 1 (currently amended) A compound corresponding to the following general formula:

nitrogen-containing aromatic ring – (NR₃)_p – (CO)_n- distribution agent –
(CO)_m – (NR'₃)_q – aromatic or non-aromatic ring

wherein

n, m, p and q are identical or different and are integers 0 or 1; and

wherein

- the nitrogen-containing aromatic ring is:
 - ◊ a quinoline optionally substituted with at least
 - one group N(Ra)(Rb) in which Ra and Rb, are identical or different, and are independently of each other hydrogen or a C1-C4 alkyl; or
 - one C1-C4 alkyl or alkoxy;
 - ◊ a quinoline possessing a nitrogen atom in quaternary form;
 - ◊ a benzamidine; or
 - ◊ a pyridine;
- the aromatic or non-aromatic ring is:
 - ◊ a quinoline optionally substituted with at least
 - one group N(Ra)(Rb) in which Ra and Rb, are identical or different, and are independently hydrogen or a C1-C4 alkyl; or
 - one C1-C4 alkyl or alkoxy;

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

- ◇ a quinoline possessing a nitrogen atom in quaternary form;
 - ~~◇ a benzamidine;~~
 - ◇ a pyridine; or
 - ◇ a phenyl optionally substituted with halogen, C1-C4 alkoxy, cyano, carbonylamino optionally substituted with one or more C1-C4 alkyl, guanyl, C1-C4 alkylthio, amino, C1-C4 alkylamino, C1-C4 dialkylamino, nitro, C1-C4 alkyleneamino or C2-C4 alkenyleneamino; or
 - ~~◇ a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic nucleus containing 0 to 2 heteroatoms per ring provided that at least one heteroatom is present in at least one ring optionally substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~
 - R₃ and R'₃, which are identical or different, represent independently of each other hydrogen or C1-C4 alkyl;
 - the distribution agent is:
 - ◇ a triazine group optionally substituted with one or more radicals chosen from halogen, C1-C4 alkyl, and thio, oxy or amino which are themselves optionally substituted with one or more C1-C4 alkyl;
 - ◇ a 5- or 6-membered heterocyclic radical containing a sulfur, oxygen or nitrogen atom;
 - ◇ a phenyl, -NH-phenyl-NH-, -NH-phenyle-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -NH-CH₂-phenyl-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl, -phenyl-CH₂-, -CH₂-thienyl-, -thienyl-CH₂-, or -CH=CH-; or
 - ◇ a diazine group; and wherein the heterocyclic, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -NH-CH₂-phenyl-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl, -phenyl-CH₂-, -CH₂-thienyl-, -thienyl-CH₂-, -CH=CH-, and diazine are optionally substituted with the same groups as the triazine;
- or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof;

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

with the proviso that:

when the distribution agent is phenyl optionally substituted with NH₂, 2, 5-pyridyl or 2,6-pyridyl, ~~and when n, m, p and q are each 1~~ and R₃ and R'₃ are hydrogen, then the nitrogen-containing aromatic ring and the aromatic ring are not both quinoline which is unsubstituted or substituted ~~on its nitrogen atom~~ with C1-C4 alkyl; ~~and~~
~~when the distribution agent is a triazine and both p and q are 1, then both n and m are not 0.~~

- 2 (original) The compound according to claim 1 which binds the G-quadruplex structure of telomeres.
- 3 (original) The compound according to claim 1 wherein the distribution agent is chosen from the heterocyclic group, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -NH-CH₂-phenyl-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl-, -phenyl-CH₂-, -CH₂-thienyl-, -thienyl-CH₂-, -CH=CH- and diazine.
- 4 (original) The compound according to claim 1 wherein the distribution agent is chosen from the heterocyclic group, the phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl-, -CH₂-thienyl-, -CH=CH-, and diazine.
- 5 (original) The compound according to claim 1 wherein the distribution agent is chosen from the heterocyclic group, the phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -CH₂-phenyl-CH₂-, -CH=CH-, and diazine.
- 6 (original) The compound according to claim 1 wherein the distribution agent is chosen from the heterocyclic group, the phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH- and diazine.
- 7 (original) The compound according to claim 1 wherein the distribution agent is thienyl or pyridyl.
- 8 (original) The compound according to claim 1 wherein the distribution agent is chosen from thienyl, pyridyl, phenyl, -NH-phenyl-NH-,

Application Ser. No.: 10/721,210

Filing Date: November 25, 2003


Examiner: Davis, Zinna Northington

-NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH- and diazine.

9 (original) The compound according to claim 1 wherein the diazine group is a pyrimidine.

10 (canceled)

11 (currently amended) The compound according to claim 1 having the following formula (IA) :



wherein

~~n, m, p and q are identical or different and are integers 0 or 1;~~

• A represents:

◊ a 5- to 6-membered heterocyclic radical containing a sulfur, oxygen or nitrogen atom;

◊ a phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -NH-CH₂-phenyl-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl, -phenyl-CH₂-, -CH₂-thienyl-, -thienyl-CH₂- or -CH=CH-; or

◊ a diazine group; and wherein the heterocyclic, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -NH-CH₂-phenyl-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl, -phenyl-CH₂-, -CH₂-thienyl-, -thienyl-CH₂-, -CH=CH-, and diazine are optionally substituted with one or more radicals chosen from halogen, C1-C4 alkyl, and thio, oxy or amino which are themselves optionally substituted with one or more C1-C4 alkyl;

- R₃ and R'₃, which are identical or different, represent independently of each other hydrogen or C1-C4 alkyl;

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

- Ar₁ and Ar₂, which are identical or different, and are independently of each other selected from:

- a quinoline optionally substituted with at least
 - a group N(Ra)(Rb) in which Ra and Rb are identical or different, and are independently of each other hydrogen or a C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
- a quinoline possessing a nitrogen atom in quaternary form;
- ~~a benzamidine;~~
- a pyridine optionally attached at the 4-position or fused with an aryl or heteroaryl group, optionally substituted with a C1-C4 alkyl; or
- a phenyl optionally substituted with halogen, C1-C4 alkoxy, cyano, carbonylamino optionally substituted with one or more C1-C4 alkyl, guanyl, C1-C4 alkylthio, amino, C1-C4 alkylamino, C1-C4 dialkylamino, nitro, C1-C4 alkyleneamino or C2-C4 alkenyleneamino; or
- ~~a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic ring containing 0 to 2 heteroatoms per ring provided that at least one heteroatom is present in at least one ring optionally substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof;

with the proviso that:

when A is phenyl optionally substituted with NH₂, 2,5-pyridyl or 2,6-pyridyl and when ~~n, m, p and q are each 1~~ and R₃ and R₃' are hydrogen, then Ar₁ and Ar₂ are not both quinoline which is unsubstituted or substituted on its nitrogen atom with C1-C4 alkyl; ~~and~~


~~when A is a triazine, and both p and q are 1, then both n and m are not 0.~~

12 (original) The compound according to claim 11 wherein A is chosen from heterocyclic group, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-,

Application Ser. No.: 10/721,210
 Filing Date: November 25, 2003
 Examiner: Davis, Zinna Northington

-NH-CH₂-phenyl-CH₂-NH-, -NH-CH₂-phenyl-NH-, -CH₂-phenyl-CH₂-,
 -CH₂-phenyl-, -phenyl-CH₂-, -CH₂-thienyl-, -thienyl-CH₂-, -CH=CH- and
 pyrimidine.

- 13 (original) The compound according to claim 11 wherein A is chosen from heterocyclic group, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -CH₂-phenyl-CH₂-, -CH₂-phenyl-, -CH₂-thienyl-, -CH=CH- and pyrimidine.
- 14 (original) The compound according to claim 11 wherein A is chosen from heterocyclic group, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, -CH₂-phenyl-CH₂-, -CH=CH- and pyrimidine.
- 15 (original) The compound according to claim 11 wherein the diazine group which A may represent is pyrimidine.
- 16 (currently amended) The compound according to claim 1 having the following formula (I) :



wherein

~~n and m are identical or different and are integers 0 or 1;~~

- A represents:

- ◇ a 5- to 6-membered heterocyclic radical containing a sulfur, oxygen or nitrogen atom;

- ◇ a phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH- or -NH-CH₂-phenyl-CH₂-NH-; or

- ◇ a diazine group; and wherein the heterocyclic, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, and diazine are optionally substituted with one or more radicals chosen from halogen, C1-C4 alkyl, and thio, oxy

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

or amino which are themselves optionally substituted with one or more C1-C4 alkyl;

- R₃ and R₃', which are identical or different, represent independently of each other hydrogen or C1-C4 alkyl;

- Ar₁ and Ar₂, which are identical or different, and are independently of each other selected from :

- a quinoline optionally substituted with at least
 - a group N(Ra)(Rb) in which Ra and Rb are identical or different, and are independently of each other hydrogen or a C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
- a quinoline possessing a nitrogen atom in quaternary form;
- ~~a benzamidine;~~
- a pyridine optionally attached at the 4-position or fused with an aryl or heteroaryl group, optionally substituted with a C1-C4 alkyl; or
- a phenyl optionally substituted with halogen, C1-C4 alkoxy, cyano, carbonylamino optionally substituted with one or more C1-C4 alkyl, guanlyl, C1-C4 alkylthio, amino, C1-C4 alkylamino, C1-C4 dialkylamino, nitro, C1-C4 alkyleneamino or C2-C4 alkenyleneamino; or
- ~~a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic ring containing 0 to 2 heteroatoms per ring provided that at least one heteroatom is present in at least one ring optionally substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof;

with the proviso that:

when A is phenyl optionally substituted with NH₂, 2,5-pyridyl or 2,6-pyridyl and when ~~n and m are 1 and~~ R₃ and R₃' are hydrogen, then Ar₁ and Ar₂ are not both quinoline which is unsubstituted or substituted ~~on its nitrogen atom~~ with C1-C4 alkyl.

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

17 (original) The compound according to claim 16 wherein A is chosen from thienyl, pyridyl, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH- and pyrimidine.

18 (canceled)

19 (original) The compound according to claim 16 wherein Ar₁ and Ar₂ represent:

- a quinoline optionally substituted with at least
 - a group N(Ra)(Rb) in which Ra and Rb are identical or different, and are independently of each other hydrogen or C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
- a quinoline possessing a nitrogen atom in quaternary form; or
- pyridine.

20 (original) The compound according to claim 16 wherein Ar₁ and Ar₂ are chosen from the following groups : 4-amino-, 4-methylamino-, 4-dimethylamino- or 4-alkoxy-quinolyl or -quinolinium in which the quinolinium is optionally substituted with one or two methyl groups.

21 (original) The compound according to claim 16 wherein A is optionally substituted with one or more radicals chosen from halogen, C1-C4 thioalkyl, amino, C1-C4 alkylamino or C1-C4 dialkylamino.

22 (original) The compound according to claim 16 wherein A is optionally substituted with methylthio or halogen.

23 – 24 (canceled)

25 (currently amended) The compound of formula (IA) according to claim 11 wherein:

n, m, p and q are ~~identical or different and are integers 0 or 1;~~

- A represents:

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

- ◇ thienyl or pyridyl;
- ◇ phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-,
-NH-CH₂-phenyl-CH₂-NH-, -CH₂-phenyl-CH₂- or -CH=CH-; or
- ◇ pyrimidyl optionally substituted with one or more radicals chosen
from halogen or C1-C4 alkylthio;
- R₃ and R'₃, which are identical or different, represent independently of
each other hydrogen or C1-C4 alkyl;
- Ar₁ and Ar₂, which are identical or different, and are independently of
each other selected from :
 - a quinoline optionally substituted with at least
 - a group N(Ra)(Rb) in which Ra and Rb are identical or
different, and are independently of each other hydrogen or
C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
 - a quinoline possessing a nitrogen atom in quaternary form; or
 - a pyridyl; or
 - ~~• a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic
ring containing 0 to 2 heteroatoms per ring provided that at least
one heteroatom is present in at least one ring optionally
substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-
C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a
pharmaceutically acceptable salt thereof.

26 (currently amended) The compound of formula (IA) according to claim 11
wherein:

~~n and m are identical or different and are integers 0 or 1, and p and q
are 1;~~

- A represents:
 - ◇ thienyl or pyridyl;
 - ◇ phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH- or
-NH-CH₂-phenyl-CH₂-NH-; or
 - ◇ pyrimidyl optionally substituted with one or more radicals
chosen from halogen or C1-C4 alkylthio;

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

- R_3 and R'_3 , which are identical or different, represent independently of each other hydrogen or C1-C4 alkyl;
- Ar_1 and Ar_2 , which are identical or different, and are independently of each other selected from :
 - a quinoline optionally substituted with at least
 - a group $N(R_a)(R_b)$ in which R_a and R_b , which are identical or different, and are independently of each other hydrogen or C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
 - a quinoline possessing a nitrogen atom in quaternary form;
or
 - a pyridyl; or
 - ~~a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic ring containing 0 to 2 heteroatoms per ring provided that at least one heteroatom is present in at least one ring optionally substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof.

- 27 (original) The compound according to claim 26 wherein Ar_1 and Ar_2 , which are identical or different, and are independently of each other chosen from the 4-amino-, 4-methylamino-, 4-dimethylamino- or 4-alkoxy-quinolyl or -quinolinium groups in which the quinolinium is optionally substituted with one or two methyl groups.
- 28 (original) The compound according to claim 26 wherein R_3 and R'_3 represent hydrogen.
- 29 (currently amended) The compound according to claim 26 wherein :
1. Ar_1 represents :
 - a quinoline substituted with at least
 - one group $N(R_a)(R_b)$ in which R_a and R_b are identical or different, and are independently of each other hydrogen or C1-C4 alkyl; or

Application Ser. No.: 10/721,210
 Filing Date: November 25, 2003
 Examiner: Davis, Zinna Northington

- a C1-C4 alkyl or alkoxy;

- a quinoline possessing a nitrogen atom in quaternary form; and

2. Ar₂ represents

- a quinoline substituted with at least
 - one group N(Ra)(Rb) in which Ra and Rb are identical or different, and are independently of each other hydrogen or C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
- a quinoline possessing a nitrogen atom in quaternary form; or
- a pyridyl;
- ~~quinoline, benzimidazole, indole, benzothiophene, benzofuran, benzothiazole, benzoxazole, carbazole, quinazoline, quinoxaline, piperidyl, piperazinyl, morpholino, azepine and diaza-azepine, which are optionally substituted by one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof.

30 (currently amended) The compound of formula (IA) according to claim 11 chosen from :

- bis[(4-methoxy-2-methylquinolin-6-yl)-amido]-2,5-thiophenedicarboxylic acid;
- bis[(4-dimethylamino-2-methylquinolin-6-yl)-amido]-2,5-thiophenedicarboxylic acid;
- bis[(4-amino-2-methylquinolin-6-yl)-amido]-2,5-thiophenedicarboxylic acid;
- N,N'-bis(4-amino-2-methylquinolin-6-yl)isophthalamide;
- N,N'-bis(4-dimethylamino-2-methylquinolin-6-yl)terephthalamide;
- 1-(4-methoxy-2-methylquinolin-6-yl)-3-{3-[3-(4-methoxy-2-methylquinolin-6-yl)ureido]phenyl}urea;
- 1-(4-dimethylamino-2-methylquinolin-6-yl)-3-[4-[3-(4-dimethylamino-2-methylquinolin-6-yl)ureido]phenyl]urea;

Application Ser. No.: 10/721,210

Filing Date: November 25, 2003

Examiner: Davis, Zienna Northington

- N,N'-bis(4-amino-2-methyl-6-quinolyl)-2,4-diamino-6-chloro-5-methyl-sulfanylpuridine;
- bis[(4-amino-2-methyl-quinolin-6-yl)-amido]-2,5-pyridinedicarboxylic acid hydrochloride;
- bis[(4-amino-2-methylquinolin-6-yl)-amido]-2,5-pyridinedicarboxylic acid;
- N,N'-bis(4-dimethylamino-2-methylquinolin-6-yl)-but-2-enediamide;
- bis[(4-dimethylamino-2-methyl-quinolin-6-yl)-amido]-2,5-pyridinedicarboxylic acid;
- bis[(4-dimethylamino-2-methylquinolin-6-yl)-amido]-2,4-pyridinedicarboxylic acid;
- N,N'-bis(4-dimethylamino-2-methylquinolin-6-yl)-1,4-phenylenediacetamide;
- bis[(4-amino-2-methyl-quinolin-6-yl)-amido]-2,6-pyridinedicarboxylic acid hydrochloride;
- bis[(4-amino-2-methyl-quinolin-6-yl)amido]-2,6-pyridine dicarboxylic acid;
- bis[(4-dimethylamino-2-methylquinolin-6-yl)amido]-2,6-pyridinedicarboxylic acid hydrochloride; and
- bis[(4-dimethylamino-2-methylquinolin-6-yl)-amido]-2,6-pyridinedicarboxylic acid;

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof.


31 (currently amended) The compound according to claim 30 chosen from :

- bis[(4-dimethylamino-2-methylquinolin-6-yl)-amido]-2,5-thiophenedicarboxylic acid;
- N,N'-bis-(4-amino-2-methylquinolin-6-yl)isophthalamide;
- 1-(4-dimethylamino-2-methylquinolin-6-yl)-3-{4-[3-(4-dimethylamino-2-methyl-quinolin-6-yl)ureido]phenyl}urea;
- N,N'-bis(4-amino-2-methyl-6-quinolyl)-2,4-diamino-6-chloro-5-methyl-sulfanylpuridine;
- bis[(4-amino-2-methylquinolin-6-yl)-amido]-2,5-pyridinedicarboxylic acid hydrochloride;
- bis[(4-amino-2-methylquinolin-6-yl)-amido]-2,5-pyridinedicarboxylic acid;

Application Ser. No.: 10/721,210
 Filing Date: November 25, 2003
 Examiner: Davis, Zinna Northington

- bis-[(4-dimethylamino-2-methylquinolin-6-yl)-amido]-2,5-pyridinedicarboxylic acid; and
 - bis[(4-dimethylamino-2-methylquinolin-6-yl)amido]-2,4-pyridinedicarboxylic acid;
- or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof.;

32 (currently amended) A pharmaceutical composition comprising therapeutically effective amount of a compound of formula (I) in combination with a pharmaceutically acceptable carrier ;



wherein

~~n and m are identical or different and are integers 0 or 1;~~

- A represents:

- ◊ a 5- to 6-membered heterocyclic radical containing a sulfur, oxygen or nitrogen atom;

- ◊ a phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH- or -NH-CH₂-phenyl-CH₂-NH-; or

- ◊ a diazine group; and wherein the heterocyclic, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, and diazine are optionally substituted with one or more radicals chosen from halogen, C1-C4 alkyl, and thio, oxy or amino which are themselves optionally substituted with one or more C1-C4 alkyl;

- R₃ and R'₃, which are identical or different, represent independently of each other hydrogen or C1-C4 alkyl;

- Ar₁ and Ar₂, which are identical or different, and are independently of each other selected from :

- a quinoline optionally substituted with at least
 - a group N(Ra)(Rb) in which Ra and Rb are identical or

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

different, and are independently of each other hydrogen or a C1-C4 alkyl; or

- a C1-C4 alkyl or alkoxy;
- a quinoline possessing a nitrogen atom in quaternary form;
- ~~• a benzamidine;~~
- a pyridine optionally attached at the 4-position or fused with an aryl or heteroaryl group, optionally substituted with a C1-C4 alkyl; or
- a phenyl optionally substituted with halogen, C1-C4 alkoxy, cyano, carbonylamino optionally substituted with one or more C1-C4 alkyl, guanyl, C1-C4 alkylthio, amino, C1-C4 alkylamino, C1-C4 dialkylamino, nitro, C1-C4 alkyleneamino or C2-C4 alkenyleneamino; or
- ~~• a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic ring containing 0 to 2 heteroatoms per ring provided that at least one heteroatom is present in at least one ring optionally substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof;

with the proviso that:

when A is phenyl optionally substituted with NH₂, 2,5-pyridyl or 2,6-pyridyl and when R₃ and R₃' are hydrogen, then Ar₁ and Ar₂ are not both quinoline which is unsubstituted or substituted with C1-C4 alkyl.


33 (original) The composition according to claim 32 which further comprises an anticancer agent.

34 (original) The composition according to claim 33 wherein the anticancer agent is chosen from alkylating agents, platinum derivatives, antibiotic agents, antimicrotubule agents, anthracyclines, group I and II topoisomerases, fluoropyrimidines, cytidine analogues, adenosine analogues, L-asparaginase, hydroxyurea, trans-retinoic acid, suramine,

Application Ser. No.: 10/721,210
 Filing Date: November 25, 2003
 Examiner: Davis, Zinna Northington

Irinotecan, topotecan, dexrazoxane, amifostine, herceptin, oestrogenic and androgenic hormones and antivasular agents.

- 35 (original) The composition according to claim 32 used in conjunction with radiation treatment.
36. (original) The composition according to claim 33 wherein each of the components is administered simultaneously, separately or sequentially.
37. (original) The composition according to claim 35 wherein the compound and the radiation treatment are administered simultaneously, separately or sequentially.
38. (currently amended) A method of treatment of a cancer in a patient comprising administering to said patient a therapeutically effective amount of a compound of formula (I):



wherein

~~n and m are identical or different and are integers 0 or 1;~~

- A represents:

- ◊ a 5- to 6-membered heterocyclic radical containing a sulfur, oxygen or nitrogen atom;

- ◊ a phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH- or -NH-CH₂-phenyl-CH₂-NH-; or

- ◊ a diazine group; and wherein the heterocyclic, phenyl, -NH-phenyl-NH-, -NH-phenyl-CH₂-NH-, -NH-CH₂-phenyl-CH₂-NH-, and diazine are optionally substituted with one or more radicals chosen from halogen, C1-C4 alkyl, and thio, oxy or amino which are themselves optionally substituted with one or more C1-C4 alkyl;

- R₃ and R'₃, which are identical or different, represent independently of

Application Ser. No.: 10/721,210
Filing Date: November 25, 2003
Examiner: Davis, Zinna Northington

each other hydrogen or C1-C4 alkyl;

- Ar₁ and Ar₂, which are identical or different, and are independently of each other selected from :

- a quinoline optionally substituted with at least
 - a group N(Ra)(Rb) in which Ra and Rb are identical or different, and are independently of each other hydrogen or a C1-C4 alkyl; or
 - a C1-C4 alkyl or alkoxy;
- a quinoline possessing a nitrogen atom in quaternary form;
- ~~a benzamidine;~~
- a pyridine optionally attached at the 4-position or fused with an aryl or heteroaryl group, optionally substituted with a C1-C4 alkyl; or
- a phenyl optionally substituted with halogen, C1-C4 alkoxy, cyano, carbonylamino optionally substituted with one or more C1-C4 alkyl, guanyl, C1-C4 alkylthio, amino, C1-C4 alkylamino, C1-C4 dialkylamino, nitro, C1-C4 alkyleneamino or C2-C4 alkenyleneamino; ~~or~~
- ~~a mono- or bi- or tricyclic aromatic or non-aromatic heterocyclic ring containing 0 to 2 heteroatoms per ring provided that at least one heteroatom is present in at least one ring optionally substituted with one or more C1-C4 alkyl, C1-C4 alkylene or C2-C4 alkenylene;~~

or an isomer, an enantiomer, a diastereoisomer or a mixture thereof, or a pharmaceutically acceptable salt thereof;

with the proviso that:

when A is phenyl optionally substituted with NH₂, 2,5-pyridyl or 2,6-pyridyl and when R₃ and R₃' are hydrogen, then Ar₁ and Ar₂ are not both quinoline which is unsubstituted or substituted with C1-C4 alkyl.